

(RESEARCH ARTICLE)



## Determination of organochlorine pesticides residues in water, fish and sediment samples from River Tella, Gassol, Taraba State, Nigeria

Haruna Ibrahim <sup>1,\*</sup> and O N Maitera <sup>2</sup>

<sup>1</sup> Department of Science Laboratory Technology, Federal Polytechnic Bali, PMB 05 Bali, Taraba State, Nigeria.

<sup>2</sup> Department of Chemistry, Modibo Adama University, P. M. B 2076, Yola, Adamawa, Nigeria.

International Journal of Scholarly Research in Science and Technology, 2023, 02(01), 019–027

Publication history: Received on 01 January 2023; revised on 08 March 2023; accepted on 10 March 2023

Article DOI: <https://doi.org/10.56781/ijrst.2023.2.1.0014>

### Abstract

The study was aimed at determining the levels of organochlorine pesticides residues in water, sediment and fish species at River Tella to find out the extent of pesticide contamination and accumulation in the river. Total of fourteen (14) organochlorine pesticide residues were analyzed which include Delta lindane, Alpha Lindane, Beta Lindane, Gamma Lindane, Heptachlor, Aldrin, Heptachlor Epoxide, Endosulfan I, Endosulfan II, P'P-DDE, P'P-DDD, P'P-DDT and Methoxychlor. The water sample was subjected to liquid-liquid extraction method while the fish and the sediment samples were subjected to soxhlet extraction. The extracts were later analyzed for organochlorine pesticide residues using Gas-Chromatography coupled with mass spectrometer (GC-MS). The study has revealed that the level of pesticides residues in water was less than that of fish and sediment which was also below the WHO/FAO values. It was also observed that high concentrations recorded were recorded in high in Gamma Lindane (24.71), follow by Alpha Lindane (23.70) and Beta Lindane (4.50). *Mormyrus rume* follow by *Clarias gariepinus* while the *Tilapia zilli* recorded low concentration of the pesticide residues. The Aldrin recorded high concentration of 5.11 ppm in *Mormyrus rume* follow by Endosulfon II which recorded the concentration of 25.03 ppm also in *Mormyrus rume*. Endrin also recorded the concentration of 4.88 ppm in *Clarias gariepinus*. The lowest pesticide residues was recorded in p,p'-DDD (0.01 ppm), p,p'-DDT (0.03 ppm), Methoxychlor (0.01 ppm). The Heptachlor epoxide and P'P-DDE were not detected in water sample. The presence of high concentration in fish may pose a great danger when these fish are being consumed over time. Regular monitoring is therefore required to control the levels of pesticide residues in the water bodies. The measurement of the fish showed that *Clarias gariepinus* had the average length of 15.89 cm, and average weight of 179.87 g. *Mormyrus rume* had the average length of 15.18 cm, and average weight of 210.54g. *Tilapia zilli* was the least species of fish sample with a length of 12.96 cm and average weight of 96.88 g. The water quality parameters of the river were also studied, which recorded the temperature of 28.89 °C was recorded, pH value of 9.01, Conductivity value of 40.11, DO value of 6.14 and TDS of 2.35.

**Keywords:** Pesticides; Residues; Organochlorine; River Tella

### 1 Introduction

Reduction of the proliferation of pest and increase in food production, has made pesticides application in agriculture inevitable [5]. Pesticides include herbicides for destroying weeds and unwanted vegetation, insecticides for controlling insects, fungicides used for preventing the growth of molds and mildew and any other substances with the similar usage [24]. Various farming, fishing and municipal activities are taking place near the bank of river Tella year in year out could bring about water quality problems and disruption in fish [2]. Pesticides constitute one of the most hazardous groups of contaminants [35], posing potential risk to humans and other life forms [20]. Thus, deaths and chronic diseases worldwide are sometimes reported to have resulted from pesticide poisoning [28]. The occurrence of pesticides residue,

\* Corresponding author: Haruna Ibrahim

especially organochlorines (OCs) in the environment is a great worry due to their tendency for long-range transport. Also their capacity to bioaccumulate in food chain poses a threat to human health and the environment [11, 19]. Pesticides enter and pollute the environment in a number of ways, including application through agriculture, accidental spillage or through the unauthorized dumping of pesticide products or their containers [12].

Chlorinated organic pesticides are very stable in both fresh and salt water and are resistant to photo degradation [3]. They will disappear from the water with secondary mechanisms such as, absorption on sediment, biological breakdown by microflora and fauna, and absorption by fish through gills, skin and feeding. They are poorly hydrolyzed and slowly biodegrades in environment. Therefore, these compounds are persistent in food chains and are readily accumulated in animal tissues. Fish absorb these compounds directly by water or by ingesting contaminated food. In particular, Organochlorine insecticides are highly stable under different environmental conditions and persistent nature and chronic adverse effects on wildlife and humans [4].

Contamination of water bodies for example is a major concern for fish and other aquatic organisms which are the major sources of protein [14]. Accumulation of pesticides in these organisms has become a serious public health issue worldwide. Fish are used extensively for environmental monitoring because they concentrate pollutants directly from water and diet, thus enabling the assessment of transfer of pollutants through the food web [9, 13, and 21]. Fish occupy different habitats in the ecosystem and have different feeding behaviors, thereby exhibiting different profile of accumulation of contaminants such as pesticides. For example, benthic fish species are considered more prone to contamination [38, 39] as they tend to accumulate sediments bound contaminants than pelagic fish [10, 27]. When large animals feed on these contaminated organisms, the toxins are taken into their bodies moving up the food chain in increasing concentration in a process known as bio magnifications [32].

The accumulation of pesticides in the sediments is because of some physicochemical parameters such as  $P^H$  which have direct influence in the solubility of these pesticides in water [29]. Sediments also serve as an important sinks and remobilization of contaminants in aquatic system. The sediments act as secondary contamination source after water in the ecosystem and are the principal reservoirs of environmental pesticides representing a source from which residues can be released to the atmosphere, ground water and taking up by living organisms [36]. Many pesticides achieve their intended use of killing pests by disrupting the nervous system, like DDT have a wide range of potential health effects. They may cause physical irritation to the skin as well as acts as carcinogens, endocrine disruption and nervous disorder [34].

The historical background of pesticides used in agriculture dated back to the beginning of agriculture itself and it became more pronounced with time due to the increased pest population parallel with decreasing in soil fertility [23]. The used of modern pesticides in agriculture and public health is dated back to the 19<sup>th</sup> century when the German Scientists Ziegler synthesize the dichlorophenyltrichloroethane (DDT) as the first important synthetic organic pesticides in 1873. About 28 years ago, it was also estimated that each year about 500 million people were affected by water-borne or associated diseases and as many as 10 million of these die [1]. Recently WHO reports that about 80 % of all human illnesses in the developing world are caused by biological contaminations [15].

Despite the good results of using pesticides in agricultural and public health as described, their use is usually accompanied with deleterious environment and public health effect. Pesticides hold a unique position among environmental contaminants due to their high biological activity and toxicity. Although some pesticides are selective in their mode of operation, their selectivity is only limited to test animals. Thus pesticides can be described as biocides (have the ability of harming all forms of life other than target pest). Since 1950, the use of pesticides has increased.

---

## 2 Material and methods

### 2.1 Study Site

The study was carried out at River Tella, Gassol Local Government Area of Taraba State, Nigeria. Gassol local government area of Taraba State lies between latitude 8°32' N and 7°26' N of the equator and longitude 10°26' E and 46°14' E of the prime meridian (Topographic sheet, 1968). It is found in dry guinea savannah. It is amongst the largest local Government in Taraba State, with an estimated land area of 5,548 km<sup>2</sup>. Based on the results of the 2006 National Population Census, Gassol local Government had a population of about 244,749 persons (NPC, 2006). It has a tropical climate marked by two seasons; dry and rainy seasons. The rainy season starts around May and ends November occasionally, with 1300 – 1400mm rainfall annually.

## 2.2 Sample Collection

Water sample was collected in 2.5 L plastic bottles (thoroughly cleaned with soap and rinsed with acetone) and covered with screwed caps from different locations at appropriate depth and turbulent midstream positions of water bodies. The sample was preserved after collection in ice to minimize degradation of pesticide, stored at 4 °C until extraction. Sediment sample were taken from positions where a fine-texture substrate deposition takes place. The upper 2cm of the bed sediment was collected with a Teflon coated spoon, stored in aluminium containers at -20 °C in the laboratory until analysis. Fish identified and selected for this project are tilapia (*Tilapia zilli*), Catfish (*Clarias gariepinus*), and Mormyrids (*Mormyrus rume*). The fish samples were collected from using standard procedures as described by [30]. Samples were collected randomly from fishermen catches at the landing sites between 9:00am – 11:00am. The measurements of the total length (cm) of each fish was taking from the tip of snout (mouth close) to the end of caudal fin using meter rule, body weight (g) was measured using electronic digital balance and the condition factor of individual fish sampled were recorded. The samples were then dissected and their organs (kidney, gills and muscles) were separated. The organs were oven dried and pounded into fine powder; and kept in a plastic container for analysis [6].

## 2.3 Sample Preparation and Extraction

Water samples was shaken well and filtered through Whatman filter paper No.1 to remove suspended materials. Sediment sample was air dried and then sieved through a 250 µm stainless steel mechanical shaker. Fish samples were thawed, cleaned with distilled water and scales sloughed off. Muscle tissues was dissected, minced into smaller pieces and homogenized. Extraction of pesticide residues in fish, water and sediment samples was done according to the method developed by [25, 30] with some modifications.

## 2.4 Extraction of pesticide residue in water sample

According to the method by [25], About 50 ml n-hexane was introduced into a 1 L separating funnel containing 100 ml of filtered water sample. The mixture was shaken vigorously for 5 min and allowed to settle. After complete separation, the organic phase was be drained into a 250 ml conical flask while the aqueous phase was re-extracted twice with 50 ml of n-hexane. The extracted organic phase was combined and dried by passing through a glass funnel packed with activated anhydrous sodium sulfate. The organic fraction was then concentrated to near dryness using vacuum rotary evaporator at 40 °C.

## 2.5 Extraction of pesticide residues in sediments

According to the method by [25], ten (10.0) g dry sediment sample was transferred into an extraction thimble. Pesticide residues in sediments were extracted with 150 ml of n-hexane/acetone mixture 4:1 v/v for 6 h using soxhlet extraction. The extract was then concentrated to near dryness using vacuum rotary evaporator at 40 °C. Each extract was dissolved in 10 ml of n-hexane and subjected to clean-up.

## 2.6 Extraction of pesticide residues in fish

According to the method by [25], Ten (10.0) g of homogenized fish sample was placed into a 100 ml conical flask and 20.0 g of activated anhydrous sodium sulfate was added and mixed thoroughly. Then 30 ml of 2:1 (v/v) hexane/acetone mixture was added and thoroughly mixed by shaking. The sample was then sonicated for 20 min using Bransonic Ultrasound Sonicator. The supernatant was filtered into a 250 ml round bottom flask. The extraction was repeated twice and all the supernatants combined and concentrated at 40 °C to near dryness using a Vacuum Rotary Evaporator.

## 2.7 Sample clean-up

Sample clean-up Approximately 2.5 g of activated silica gel was weighed and parked into a glass column which has been plugged with glass wool and 1.0 g of anhydrous sodium sulfate. About 10 ml n-hexane was used to wet and rinse the column. The extract was then transferred into the column and eluted with 20 mL portions of hexane/acetone mixtures. The eluates was collected into a round bottomed flask and then concentrated to dryness. The residues were then be dissolved in 2 mL of ethyl acetate and placed in a GC vial for gas chromatograph analysis.

## 2.8 GC Analysis

The GC analysis of the organochlorine pesticide residues was conducted using a model 2010 Shimadzu GC equipped with an EC. Separation was done on an SGE BPX-5 of 60 m capillary column with 0.25 mm internal diameter and 0.25 µm film thicknesses, equipped with 1 m retention gap. The oven temperature was programmed as follows: initial temperature was set at 90 °C for 3 min and ramped at 30 °C/min to 200 °C for 15 min and then to 265 °C at a rate of 5 °C/min for 5 min then to 275 °C at the rate of 3 °C/min and allowed to stay for 15 min. The injector setting is a pulsed

splitless mode with a temperature of 250 °C at a pressure of 1.441 bar. Pulsed pressure was 4.5 bar, pulsed time 1.5 min, purge flow of 55.4 mL/min with a purge time of 1.4 min. The detector temperature was 300 °C. Nitrogen was used as carrier gas at a flow rate of 30 mL/min. A Varian CP-3800 GC equipped with a Combi PAL Auto sampler was used to measure levels of the pesticide residues. The column used was a 30 m × 0.25 mm internal diameter fused silica capillary coated with VF-1701 (0.25 µm film). The oven temperature was programmed as follows: initial temperature was set at 65 °C for 2 min and ramped at 25 °C/min to 210 °C for 6 min and then to 230 °C at 20 °C/min and allowed to stay for 20 min. The injector setting is a pulsed splitless mode at a temperature of 230 °C. The detector temperature was 250 °C in “constant make-up flow” mode. Helium gas was used as carrier gas at a flow rate of 2 mL/min.

### 3 Results and discussion

The tables below are the results of the organochlorine pesticide residues analysis in water, sediments and the three fish species samples collected from River Tella. The results are measured in concentration (ppm).

**Table 1** Concentration (ppm) of pesticide residues in water samples at river Tella

Pesticides residues	Types of residue	Molecular Formula	R.T. (min)	Concentration (ppm)
Delta.-Lindane	Insecticide	C <sub>6</sub> H <sub>6</sub> Cl <sub>6</sub>	5.353	0.03
Alpha.-Lindane	Insecticide	C <sub>6</sub> H <sub>6</sub> Cl	5.839	0.09
Beta.-Lindane	Insecticide	C <sub>6</sub> H <sub>6</sub> Cl <sub>6</sub>	6.040	0.25
Gamma.-Lindane	Insecticide	C <sub>6</sub> H <sub>6</sub> Cl <sub>6</sub>	6.549	0.04
Heptachlor	Insecticide	C <sub>10</sub> H <sub>5</sub> Cl <sub>7</sub>	6.904	0.01
Aldrin	Herbicide	C <sub>12</sub> H <sub>8</sub> Cl <sub>6</sub>	7.327	0.50
Heptachlor epoxide	Insecticide	C <sub>10</sub> H <sub>5</sub> Cl <sub>7</sub>	0.000	N.D.
Endosulfan I	Insecticide	C <sub>9</sub> H <sub>6</sub> Cl <sub>6</sub> O <sub>3</sub> S	8.626	0.06
p,p'-DDE	Insecticide	C <sub>14</sub> H <sub>8</sub> Cl <sub>4</sub>	9.101	0.00
Endrin	Insecticide	C <sub>12</sub> H <sub>8</sub> Cl <sub>6</sub> O	9.444	0.02
Endosulfan II	Insecticide	C <sub>9</sub> H <sub>6</sub> Cl <sub>6</sub> O <sub>3</sub> S	9.770	0.17
p,p'-DDD	Insecticide	C <sub>14</sub> H <sub>10</sub> Cl <sub>4</sub>	9.925	0.03
p,p'-DDT	Insecticide	C <sub>14</sub> H <sub>9</sub> Cl <sub>5</sub>	10.491	0.03
Methoxychlor	Insecticide	C <sub>6</sub> H <sub>15</sub> Cl <sub>3</sub> O <sub>2</sub>	11.498	0.01

The morphometric characteristics such as the average length, average weight and condition factor of the studied fish species were measured. The *Clarias gariepinus* had the highest average length of 15.89 cm, and average weight of 179.87 g. *Mormyrus rume* had the medium average length of 15.18 cm, and average weight of 210.54 g. *Tilapia zilli* was the least species of fish sample with a length of 12.96 cm and average weight of 96.88 g. The *Tilapia zilli* had highest condition factor of 4.45, follow by *C. gariepinus* which recorded the K- factor of 4.41 while the *Mormyrus rume* had the least K-factor of 3.86. The condition factor (K) gives information on the physiological condition of fish in relation to its welfare. [26] Reported that fishes with a low condition index are presumably believed to have experienced adverse physical environment or insufficient nutrition. [7] Recommended K value range (2.9-4.8) as suitable for matured fresh water fish. It was observed that the three fish species sampled in this study recorded varying condition factors, all the samples shows slight increases in K - value. Also, all of the fish species sampled in this present study had condition factors > 1, and were within the normal ranges as recommended by [31] who stated that condition factor greater or equal to one is good, indicating a good level of feeding, and proper environmental condition

The water quality parameters of the river were also studied. The temperature of 28.89 °C was recorded, pH value of 9.01, Conductivity value of 40.11, DO value of 6.14 and TDS of 2.35. Water quality testing is an important part of environmental monitoring. When water quality is poor, it affects not only aquatic life but the surrounding ecosystem as well.

**Table 2** Concentration (ppm) of pesticide residues in *Tilapia zilli* from Tella River

Pesticides residues	Types of residue	Molecular Formula	R.T. (min)	Concentration (ppm)
Delta.-Lindane	Insecticide	C <sub>6</sub> H <sub>6</sub> Cl <sub>6</sub>	5.364	0.22
Alpha.-Lindane	Insecticide	C <sub>6</sub> H <sub>6</sub> Cl	5.802	0.45
Beta.-Lindane	Insecticide	C <sub>6</sub> H <sub>6</sub> Cl <sub>6</sub>	6.051	2.36
Gamma.-Lindane	Insecticide	C <sub>6</sub> H <sub>6</sub> Cl <sub>6</sub>	6.554	0.32
Heptachlor	Insecticide	C <sub>10</sub> H <sub>5</sub> Cl <sub>7</sub>	6.755	0.11
Aldrin	Herbicide	C <sub>12</sub> H <sub>8</sub> Cl <sub>6</sub>	7.310	0.60
Heptachlor epoxide	Insecticide	C <sub>10</sub> H <sub>5</sub> Cl <sub>7</sub>	8.014	0.14
Endosulfan I	Insecticide	C <sub>9</sub> H <sub>6</sub> Cl <sub>6</sub> O <sub>3</sub> S	8.620	0.36
p,p'-DDE	Insecticide	C <sub>14</sub> H <sub>8</sub> Cl <sub>4</sub>	9.072	0.09
Endrin	Insecticide	C <sub>12</sub> H <sub>8</sub> Cl <sub>6</sub> O	9.478	0.24
Endosulfan II	Insecticide	C <sub>9</sub> H <sub>6</sub> Cl <sub>6</sub> O <sub>3</sub> S	9.782	0.96
p,p'-DDD	Insecticide	C <sub>14</sub> H <sub>10</sub> Cl <sub>4</sub>	9.907	0.01
p,p'-DDT	Insecticide	C <sub>14</sub> H <sub>9</sub> Cl <sub>5</sub>	10.491	0.12
Methoxychlor	Insecticide	C <sub>6</sub> H <sub>15</sub> Cl <sub>3</sub> O <sub>2</sub>	11.481	0.05

**Table 3** Concentration (ppm) of pesticide residues in *Clarias gariepinus* from Tella River

Pesticides residues	Types of residue	Molecular Formula	R.T. (min)	Concentration (ppm)
Delta.-Lindane	Insecticide	C <sub>6</sub> H <sub>6</sub> Cl <sub>6</sub>	5.335	0.30
Alpha.-Lindane	Insecticide	C <sub>6</sub> H <sub>6</sub> Cl	5.850	0.07
Beta.-Lindane	Insecticide	C <sub>6</sub> H <sub>6</sub> Cl <sub>6</sub>	6.045	1.22
Gamma.-Lindane	Insecticide	C <sub>6</sub> H <sub>6</sub> Cl <sub>6</sub>	6.554	0.62
Heptachlor	Insecticide	C <sub>10</sub> H <sub>5</sub> Cl <sub>7</sub>	6.749	0.07
Aldrin	Herbicide	C <sub>12</sub> H <sub>8</sub> Cl <sub>6</sub>	7.344	1.04
Heptachlor epoxide	Insecticide	C <sub>10</sub> H <sub>5</sub> Cl <sub>7</sub>	7.996	0.7
Endosulfan I	Insecticide	C <sub>9</sub> H <sub>6</sub> Cl <sub>6</sub> O <sub>3</sub> S	8.626	0.75
p,p'-DDE	Insecticide	C <sub>14</sub> H <sub>8</sub> Cl <sub>4</sub>	9.043	0.04
Endrin	Insecticide	C <sub>12</sub> H <sub>8</sub> Cl <sub>6</sub> O	9.455	2.80
Endosulfan II	Insecticide	C <sub>9</sub> H <sub>6</sub> Cl <sub>6</sub> O <sub>3</sub> S	9.764	4.88
p,p'-DDD	Insecticide	C <sub>14</sub> H <sub>10</sub> Cl <sub>4</sub>	9.919	0.18
p,p'-DDT	Insecticide	C <sub>14</sub> H <sub>9</sub> Cl <sub>5</sub>	10.502	0.36
Methoxychlor	Insecticide	C <sub>6</sub> H <sub>15</sub> Cl <sub>3</sub> O <sub>2</sub>	11.498	0.13

**Table 4** Concentration (ppm) of pesticide residues in *Mormyrus rume* from Tella river

Pesticides residues	Types of residue	Molecular Formula	R.T. (min)	Concentration (ppm)
Delta.-Lindane	Insecticide	C <sub>6</sub> H <sub>6</sub> Cl <sub>6</sub>	5.370	0.12
Alpha.-Lindane	Insecticide	C <sub>6</sub> H <sub>6</sub> Cl	5.839	0.15
Beta.-Lindane	Insecticide	C <sub>6</sub> H <sub>6</sub> Cl <sub>6</sub>	6.051	0.07
Gamma.-Lindane	Insecticide	C <sub>6</sub> H <sub>6</sub> Cl <sub>6</sub>	6.566	0.11
Heptachlor	Insecticide	C <sub>10</sub> H <sub>5</sub> Cl <sub>7</sub>	6.766	0.04
Aldrin	Herbicide	C <sub>12</sub> H <sub>8</sub> Cl <sub>6</sub>	7.356	5.11
Heptachlor epoxide	Insecticide	C <sub>10</sub> H <sub>5</sub> Cl <sub>7</sub>	8.013	0.01
Endosulfan I	Insecticide	C <sub>9</sub> H <sub>6</sub> Cl <sub>6</sub> O <sub>3</sub> S	8.643	0.50
p,p'-DDE	Insecticide	C <sub>14</sub> H <sub>8</sub> Cl <sub>4</sub>	9.078	0.10
Endrin	Insecticide	C <sub>12</sub> H <sub>8</sub> Cl <sub>6</sub> O	9.444	2.42
Endosulfan II	Insecticide	C <sub>9</sub> H <sub>6</sub> Cl <sub>6</sub> O <sub>3</sub> S	9.781	5.03
p,p'-DDD	Insecticide	C <sub>14</sub> H <sub>10</sub> Cl <sub>4</sub>	9.896	0.09
p,p'-DDT	Insecticide	C <sub>14</sub> H <sub>9</sub> Cl <sub>5</sub>	10.480	0.09
Methoxychlor	Insecticide	C <sub>6</sub> H <sub>15</sub> Cl <sub>3</sub> O <sub>2</sub>	11.509	0.13

**Table 5** Concentration (ppm) of pesticide residues in sediment from Tella river

Pesticides residues	Types of residue	Molecular Formula	R.T. (min)	Concentration (ppm)
Delta.-Lindane	Insecticide	C <sub>6</sub> H <sub>6</sub> Cl <sub>6</sub>	5.359	2.68
Alpha.-Lindane	Insecticide	C <sub>6</sub> H <sub>6</sub> Cl	5.908	23.70
Beta.-Lindane	Insecticide	C <sub>6</sub> H <sub>6</sub> Cl <sub>6</sub>	6.051	4.50
Gamma.-Lindane	Insecticide	C <sub>6</sub> H <sub>6</sub> Cl <sub>6</sub>	6.532	24.71
Heptachlor	Insecticide	C <sub>10</sub> H <sub>5</sub> Cl <sub>7</sub>	6.772	0.18
Aldrin	Herbicide	C <sub>12</sub> H <sub>8</sub> Cl <sub>6</sub>	7.316	1.80
Heptachlor epoxide	Insecticide	C <sub>10</sub> H <sub>5</sub> Cl <sub>7</sub>	8.036	0.19
Endosulfan I	Insecticide	C <sub>9</sub> H <sub>6</sub> Cl <sub>6</sub> O <sub>3</sub> S	8.626	2.12
p,p'-DDE	Insecticide	C <sub>14</sub> H <sub>8</sub> Cl <sub>4</sub>	9.078	0.08
Endrin	Insecticide	C <sub>12</sub> H <sub>8</sub> Cl <sub>6</sub> O	9.467	1.89
Endosulfan II	Insecticide	C <sub>9</sub> H <sub>6</sub> Cl <sub>6</sub> O <sub>3</sub> S	9.793	2.28
p,p'-DDD	Insecticide	C <sub>14</sub> H <sub>10</sub> Cl <sub>4</sub>	9.902	0.22
p,p'-DDT	Insecticide	C <sub>14</sub> H <sub>9</sub> Cl <sub>5</sub>	10.498	0.36
Methoxychlor	Insecticide	C <sub>6</sub> H <sub>15</sub> Cl <sub>3</sub> O <sub>2</sub>	11.470	0.17

A total of fourteen (14) organochlorine pesticide residues were analyzed which include Delta lindane, Alpha Lindane, Beta Lindane, Gamma Lindane, Heptachlor, Aldrin, Heptachlor Epoxide, Endosulfan I, Endosulfan II, P'P-DDE, P'P-DDD, P'P-DDT and Methoxychlor. Based on the result of the analysis of the pesticide residue on the water samples collected as indicated in table 1 above, the Aldrin (0.50 ppm) was found to be higher in all the water samples, follow by Beta.-

Lindane (0.25 ppm) and Endosulfan II (0.17 ppm). The lowest pesticide residues were recorded in p,p'-DDD, p,p'-DDT and Methoxychlor. The Heptachlor epoxide and p,p'-DDE were not detected in the water sample collected. The result also indicated that the levels of pesticide residues in water are generally lower than that of fish and sediment because these pesticides are lipophilic and are not soluble in water. Also, the concentration of pesticide residue in water was below maximum residue limit (MRL) as compared with [15, 16] where the pesticide residue limit ranges from 0.001 to 0.5 ppm.

The result on the fish analysis shows that the concentrations of pesticides residues in fish were found to be higher than that of the water sample. Also, almost all the fourteen (14) organochlorine pesticides analyzed were found to be present in the fish samples collected. Based on the result obtained, it was also observed that these concentrations recorded high in *Mormyrus rume* followed by *C. gariepinus*, while the *Tilapia zilli* recorded low concentration of the pesticide residues. The Aldrin recorded high concentration of 5.11 ppm in *Mormyrus rume* followed by Endosulfan II which recorded the concentration of 5.03 ppm also in *Mormyrus rume* and 4.88 ppm in *C. gariepinus* as shown in tables 3 & 4 above. The lowest pesticide residues were recorded in p,p'-DDD, p,p'-DDE, p,p'-DDD, Methoxychlor and Heptachlor epoxide. Also, the high concentration of pesticide observed in *Clarias gariepinus* and *Mormyrus rume* may be attributed to the feeding mode of the fish [18]. This result is corroborated by [8] who related to habitation and feeding habit of *Clarias gariepinus*. To an increased concentration of pesticide residues compared with other fish species. [33], equally adds that pesticides accumulation in fish was due to their lipid content; this implies that due to the high lipid content in *C. gariepinus*, and *Mormyrus rume* more pesticide residues tend to be trapped in their lipid stores.

The level of pesticide residues in sediment analyzed was higher than that of both the water and fish samples. Also, all the fourteen (14) organochlorine pesticides analyzed were detected in the sediment sample collected as indicated in table 5 above. Gamma-Lindane recorded high concentration of 24.71 ppm followed by Alpha-Lindane which recorded the concentration of 23.70 ppm and Beta-Lindane 4.50 ppm. The lowest pesticide residue was recorded in p,p'-DDE, p,p'-DDD, p,p'-DDT, Methoxychlor and Heptachlor epoxide which were also below the MRL by [16, 17].

The low level of p,p'-DDE, p,p'-DDD, p,p'-DDT, Methoxychlor and Heptachlor epoxide in the water sample, fish and sediment showed that the farmers around the lakes and rivers do not use them much in their farming activities. However, these organochlorine pesticides were measured high in fish and sediment samples, which is because they are less soluble in water. They, therefore, accumulate in fishes and sediments when they are discharged into water bodies [37]. The lake sediments act as a sink for the persistent contaminants, whose resuspension during the lake's mixing may increase pesticide bioavailability and accumulation in the fish [18]. Pesticide pollution to the lake is therefore, likely to pose a danger to both aquatic organisms and humans [22]

---

#### 4 Conclusion

This study has revealed levels of accumulation and contamination on the different pesticide residues in water, fish and sediment samples collected from River Tella, Gassol, Taraba State. The concentration of pesticide levels in water which was less than that of fish and sediment which was also below the WHO/FAO values. It was also observed that these concentrations recorded high in *Mormyrus rume*, followed by *Clarias gariepinus*, while the *Tilapia zilli* recorded low concentration of the pesticide residues. It is recorded that the concentration of these pesticides residues were very high in fish samples. The high levels of these pesticides may pose a great danger when these fish are being consumed over time. Regular monitoring is therefore required to control the levels of pesticide residues in the water bodies.

---

#### Compliance with ethical standards

##### *Acknowledgments*

The authors would like to thank the technologist at the ABTI American University for providing the laboratory equipment for the analysis and also the field assistance for the samples collection.

##### *Disclosure of conflict of interest*

The authors declare that they have no conflict of interest.

---

**References**

- [1] Abera, S. (2011). Bacteriological analyses of drinking water sources, Department of Medical Laboratory Science and Pathology, Jima University Ethiopia. *African Journal of microbiology research*. Vol 5 (18). Online via <http://www.academicjournals.org/ajmr>.
- [2] Abu-Hilal AH (1993) Observations on heavy metal geochemical association in marine Sediments of the Jordan Gulf of Aqaba. *Mar Pollut Bull* 26: 85-90.
- [3] Abu-Hilal AH, Badran MI (1990) Effect of pollution sources on metal concentration in sediment cores from the Gulf of Aqaba (Red Sea). *Mar Pollut Bull* 21: 190-197.
- [4] Abu-Hilal AH (1987) Distribution of trace elements in nearshore surface sediments from the Jordan Gulf of Aqaba (Red Sea). *Mar Pollut Bull* 18: 190-193
- [5] Akoto O, Andoh H, Darko G, Eshun K, Osei-Fosu P (2013). Health risk assessment of pesticides residue in maize and cowpea from Ejura, Ghana. *Chemosphere* 92:67–73.
- [6] Ayelaja, A. A., George, F. O. A., Shorinmade, A. Y., Jimoh, W. A., Afolabi, Q. O. and Olawepo, K. D. (2014). Heavy Metal Concentration in Selected fish Species from Eleyele Reservoir Ibadan, Oyo State, South-West Nigeria. *Africa Journ. of Environmental Science and tech.*, 8(7): 422 – 427.
- [7] Bagenal T.B. and Tesch F.W. (1978). *Methods of Assessment of Fish Production in Fresh Waters*. IBP Handbook No. 3, 3rd ed. Oxford Blackwell Scientific Publication, London. pp. 101-136
- [8] Biego GHM, Yao KD, Ezoua P, KouadioLP, (2010). Assessment of Organochlorine Pesticides Residues in fish sold in Abidjan Markets and fishing sites. *At. J. food Agric. Nutr. Dev*, 10:3
- [9] Bruggeman WA (1982) Bioaccumulation and transformation of dichlorobiphenyls in fish. In: Hutzinger O (ed) *The handbook of environmental chemistry*, vol 2, Part B, pp 29–47
- [10] Ccancapa A, Masiá A, Navarro-Ortega A, Picó Y, Barceló D (2016), Pesticides in the Ebro River basin: occurrence and risk assessment. *Environ Pollution*, 211:414–424.
- [11] Chau KW (2005), Characterization of transboundary POP contamination in aquatic ecosystems of Pearl River delta. *Mar Pollut Bull* 51:960–965.
- [12] Chau KW (2005), Characterization of transboundary POP contamination in aquatic ecosystems of Pearl River delta. *Mar Pollut Bull* 51:960–965.
- [13] Cox JR (2002), Pesticide residue analysis facilities: experiences from the Natural Resource Institute's Support Program. In: Hanak E, Boutrif E, Fabre P, Pineiro M (Scientific Editors) *Food safety management in developing countries*, Proceedings International Workshop, CIRAD-FAO, December 11–13, 2000. Montpellier, France.
- [14] Das B, Khan YSA, Das P, Shaheen SM (2002), Organochlorine pesticide residues in catfish, *Tachysurus thalassinus* (Ruppell, 1835) from the South Patches of the Bay of Bengal. *Environ Pollut* 120:255–259.
- [15] Essumang DK, Chokky L (2009), Pesticide residues in the water and fish (Lagoon tilapia) samples from Lagoons in Lagoons in Ghana. *Bull Chem Soc Ethiopia* 23(1):19–27
- [16] FAO/WHO (2009) Food standard programme. In: *Codex Alimentarius Commission*, vol 9 (4), pp 149–158
- [17] FAO/WHO (2010) Pesticide residues in food and feed. Acceptable daily intake; *Codex Alimentarius Commission*. FAO/WHO Food Standards, Rome
- [18] Fianko R. J, Augustine D, Samuel TL, Paul OY, Eric TG, Theodosia A, Augustine F (2011), Health Risk Associated with pesticide Contamination of fish from the Densu River Basin in Ghana, *J. Environ. Protection* 2:115-123.
- [19] Getenga ZM, Kengara FO, Wandiga SO (2004) Determination of organochlorine pesticides in soil and water from river Nyando drainage system within Lake Victoria Basin, Kenya. *Bull Environ Contam Toxicol* 72: 335-343.
- [20] Guo Y, Meng X, Tang H, Zeng EY (2008), Tissue distribution of organochlorine pesticides in fish collected from the Pearl River Delta, China: implications for fishery input source and bioaccumulation. *Environ Pollut* 155:150–156.
- [21] Jeyakumar T, Kalaiarasi I, Rajavel AL, Anbu M, Kumar R (2014), Levels of organochlorine pesticide residues in water and sediment from selected agricultural sectors of Kanyakumari District, Tamil Nadu, *India. Int J Environ Res* 8(2):493–500.



- [22] Lanfranchi AL, Menone ML, Miglioranza KSB, Janiot LJ *et al* (2006), Striped weakfish (*Cynoscion guatucupa*): a biomonitor of organochlorine pesticides in estuarine and near-coastal zones. *Mar Pollut Bull* 52:74–80.
- [23] Mavura WJ, Wangila PT (2014) Distribution of Pesticide Residues in Various Lake Matrices: Water, Sediment, Fish And Algae, The Case Of Lake Nakuru, Kenya. The African network for chemical analysis of pesticides, Arusha
- [24] Muir P. (2002). The history of pesticide use, Oregon State University Press, US.
- [25] Neil, P. (2015). Environmental Chemistry, (3rd Edition) Chapman and Hall, London pp68- 200.
- [26] Okoya AA, Ogunfowokan AO, Asubiolo OI, Torto N (2013). Organochlorine Pesticide Residues in Sediments and Waters from Cocoa Producing Areas of Ondo State, Southwestern Nigeria. *International Scholarly Research Network* (ISRN) Soil Science.
- [27] Perry R.I., Hargreaves N.B., Waddell B.J. and Mackas L. (1996). Spatial variations in feeding and condition of juvenile pink and chum salmon off Vancouver Island, British Columbia. *Fish Oceanogr.*, 5 (2), 73-88
- [28] Qadir A, Malik RN (2011), Heavy metals in eight edible fish species from two polluted tributaries (Aik and Palkhu) of the River Chenab. *Biol Trace Elem Res, Pakistan*. doi:10.1007/s12011-011-9011-3.
- [29] Rigotto RM, da Silva AMC, Ferreira MJM *et al.* (2013), Trends of chronic health effects associated to pesticide use in fruit farming regions in the state of Ceará, Brazil. *Rev Bras Epidemiol* 16(3):763–773.
- [30] Swati, S. (2015). Analyses of pesticides in water, sediment and fish tissue from river Deomoni flowing through the teagardens of terai Region of west Bengal, *India International Journal of fisheries and aquatic studies* 3 (2); 17-23.
- [31] Therdteppitak A, Yammang K (2002) Determination of organochlorine pesticides in commercial fish by gas chromatography with electron capture detector and confirmation by gas chromatography–mass spectrometry. *Sci Asia* 29(2003):127–134 doi:10.12691/aees-1-3-2.
- [32] Ujjania N.C., Kohli M.P.S. and Sharma L.L. (2012). Length-weight relationship and condition factors of Indian major carps (*C. catla*, *L. rohita* and *C. mrigala*) in Mahi Bajaj Sagar, *India. Research Journal of Biology*, 2 (1), 30-36
- [33] UNEPA (2007), Volunteer Lake Monitoring, A methods Manual. Office of water US Environmental Protection Agency, Washinton DC.
- [34] Upadhi F, Wokoma QAF (2012), Examination of some pesticides residues in surface water, sediment and fish tissue of Eiechi Creak, Niger Delta, Nigeria Res. *Eviron., Earth Sci.* 4(11):939-944.
- [35] USEPA (2000), Guidance for Assessing Chemical Contamination Data for Use in fish Advisories; EPA 823-B-00-007. (3rd Edition), Vol. 1 office of Science and Tech. and Office of Water, USEPA Washinton DC, USA.
- [36] Vega AB, Frenich AG, Vidal JLM (2005), Monitoring of pesticides in agricultural water and soil samples from Andalusia by liquid chromatography coupled to mass spectrometry. *Anal Chim Acta* 538:117–127.
- [37] Walley, F. Tailoy and Lupwagi, P. (2006). Herbicides effects on pulse crop modulation and Nitrogen fixation Teach 2006 proceeding pp, 121-123.